



**Figure 3.2-28.** Historical Red King Crab Catch (in 1,000 mt) in Cook Inlet, Kodiak, Aleutian Islands, and Bristol Bay. Cook Inlet, Kodiak, and Aleutian Islands catches are given per season (for example, 1960 refers to the 1960/61 season).




Source: ADF&G 2000b

**Figure 3.2-29.** Historical Blue King Crab Catch (in 1,000 mt) in the Pribilof Islands and St. Matthew Island. The Pribilof Islands catches are given per season (for example, 1972 refers to the 1972/73 season).




Source: ADF&G 2000b

**Figure 3.2-30.** Historical Seasonal Golden King Crab Catch (in 1,000 mt) in PWS and the Aleutian Islands. Catches are given per season (for example, 1981 refers to the 1981/82 season).




Source: ADF&G 2000b

**Figure 3.2-31.** Historical Tanner Crab Catch (in 1,000 mt) in the PWS, Cook Inlet, Kodiak, Chignik, Southern Peninsula, and Bering Sea Areas. Some catches are given per season (spanning two calendar years), whereas others are per calendar year.




Source: ADF&G 2000b

**Figure 3.2-32.** Historical Snow Crab Catch (in 1,000 mt) in the Bering Sea. Some catches are given per season (spanning two calendar years), whereas others are per calendar year.




Source: ADF&G 2000b

**Figure 3.2-33.** Historical Korean Hair Crab Catch (in 1,000 mt) in the Bering Sea. Some catches are given per season (spanning two calendar years), whereas others are per calendar year.




Source: ADF&G 2000b

**Figure 3.2-34.** Historical Annual Dungeness Crab Catch (in 1,000 mt) in the PWS, Cook Inlet, Kodiak, and Alaska Peninsula Districts.




Source: ADF&G 2000b

**Figure 3.2-35.** Management Areas Involving Prohibited Species and Walrus in the Bering Sea




Source: NMFS 2001a

**Figure 3.2-36.** Historical Trend of Pacific Herring Catch in the Bering Sea and the Gulf of Alaska, 1990-1998.




Source: ADF&G Fish Ticket Landing Receipt Database, 1999

Figure 3.2-37. Final Critical Habitat for Spectacled and Steller's Eider




Source: NMFS 2001b

**Figure 3.2-38.** Distribution of Horny Corals (kilograms per hectare [kg/ha])




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

**Figure 3.2-39.** Distribution of Sponges (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

**Figure 3.2-40.** Distribution of Bryozoans (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

**Figure 3.2-41.** Distribution of Hydroids (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-42. Distribution of Sea Raspberries (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

**Figure 3.2-43.** Distribution of Sea Pen (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

**Figure 3.2-44.** Distribution of Anemones (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-45. Distribution of Sea Onion (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

Figure 3.2-46. Distribution of Sea Peach (kg/ha)




Source: <http://www.afsc.noaa.gov/groundfish/HAPC/EBScontents.htm>

**Figure 3.3-1. Biomass Trends in Bering Sea Trophic Guilds, 1979 to 1998**




Source: NMFS 2001b

Figure 3.3-2. Trophic Interactions of Key Eastern Bering Sea Groundfish




Source: NMFS 2001a

Figure 3.3-3. Trophic Relationships of the Groundfishes in the Aleutian Islands




Source: NMFS 2001a

**Figure 3.3-4.** Trophic Relationships of the Groundfishes in the Gulf of Alaska




Source: NMFS 2001a

**Figure 3.3-5.** Trend in Index of Species Composition Based on Ordination of Species Abundance Data from Five Triennial Surveys on Gulf of Alaska Shelf and Slope with Approximate 95 Percent Confidence Interval



Source: NMFS 2001a

**Figure 3.4-1. Pelagic Trawl**




Source: Jennings et al. 2001  
Permission to use figure requested by Craig Rose (pending)

**Figure 3.4-2. Bottom Trawl**




Source: K. Richards, Tetra Tech FW, Bothell, WA

**Figure 3.4-3. Set Longline Gear**




Source: K. Richards, Tetra Tech FW, Bothell, WA

**Figure 3.4-4.** Vertical Longline




Source: K. Richards, Tetra Tech FW, Bothell, WA

**Figure 3.4-5.** Jig Machine



Source: K. Richards, Tetra Tech FW, Bothell, WA

Figure 3.4-6. Gulf of Alaska: 1990-2002 Observed Pollock Pelagic Trawl Effort (Sets/25 km<sup>2</sup>)



Note: (Sets/25 km<sup>2</sup>) = Summed for 1990 to 2002  
Source: Cathy Coon, NPFMC Staff, North Pacific Observer Program Data